

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 9/20, 9/48, 31/415		A1	(11) International Publication Number: WO 00/47188
			(43) International Publication Date: 17 August 2000 (17.08.00)
(21) International Application Number: PCT/US00/03899			(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 11 February 2000 (11.02.00)			
(30) Priority Data: 60/119,638 11 February 1999 (11.02.99) US			
(71) Applicant (for all designated States except US): EMISPHERE TECHNOLOGIES, INC. [US/US]; 765 Old Saw Mill River Road, Tarrytown, NY 10591 (US).			
(72) Inventor; and (75) Inventor/Applicant (for US only): GSCHNEIDNER, David [US/US]; 44 Ceretta Street, Stamford, CT 06907 (US).			
(74) Agents: LESSLER, Jay, P. et al.; Darby & Darby P.C., 805 Third Avenue, New York, NY 10022-7513 (US).			Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: OXADIAZOLE COMPOUNDS AND COMPOSITIONS FOR DELIVERING ACTIVE AGENTS			
(57) Abstract Oxadiazole compounds and compositions for the delivery of active agents are provided. Methods of administration and preparation are provided as well.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

5

OXADIAZOLE COMPOUNDS AND COMPOSITIONS

10

FOR DELIVERING ACTIVE AGENTS

15

20

FIELD OF THE INVENTION

The present invention relates to oxadiazole compounds for delivering active agents, such as biologically or chemically active agents, to a target. These compounds are well suited for forming non-covalent mixtures with active agents for oral, intracolonic, or other routes of administration to animals. Methods for the preparation and administration of such compositions are also disclosed.

30

BACKGROUND OF THE INVENTION

Conventional means for delivering active agents are often severely limited by biological, chemical, and physical barriers. Typically, these barriers are imposed by the environment through which delivery occurs, the environment of the target for delivery, and/or the target itself. Biologically and chemically active agents are particularly vulnerable to such barriers.

In the delivery to animals of biologically active and chemically active pharmacological and therapeutic agents, barriers are imposed by the body. Examples of physical barriers are the skin, lipid bi-layers and various organ
5 membranes that are relatively impermeable to certain active agents but must be traversed before reaching a target, such as the circulatory system. Chemical barriers include, but are not limited to, pH variations in the gastrointestinal (GI) tract and degrading enzymes.

10 These barriers are of particular significance in the design of oral delivery systems. Oral delivery of many biologically or chemically active agents would be the route of choice for administration to animals if not for biological, chemical, and physical barriers. Among the numerous agents
15 which are not typically amenable to oral administration are biologically or chemically active peptides, such as calcitonin and insulin; polysaccharides, and in particular mucopolysaccharides including, but not limited to, heparin; heparinoids; antibiotics; and other organic substances. These
20 agents may be rapidly rendered ineffective or destroyed in the gastro-intestinal tract by acid hydrolysis, enzymes, and the like. In addition, the size and structure of macromolecular drugs may prohibit absorption.

Earlier methods for orally administering vulnerable
25 pharmacological agents have relied on the co-administration of adjuvants (e.g., resorcinols and non-ionic surfactants such as polyoxyethylene oleyl ether and n-hexadecylpolyethylene ether) to increase artificially the permeability of the intestinal walls, as well as the co-administration of enzymatic
30 inhibitors (e.g., pancreatic trypsin inhibitors, diisopropylfluorophosphate (DFF) and trasylol) to inhibit


enzymatic degradation. Liposomes have also been described as drug delivery systems for insulin and heparin. However, broad spectrum use of such drug delivery systems is precluded because: (1) the systems require toxic amounts of adjuvants or
5 inhibitors; (2) suitable low molecular weight cargos, i.e. active agents, are not available; (3) the systems exhibit poor stability and inadequate shelf life; (4) the systems are difficult to manufacture; (5) the systems fail to protect the active agent (cargo); (6) the systems adversely alter the
10 active agent; or (7) the systems fail to allow or promote absorption of the active agent.

More recently, proteinoid microspheres have been used to deliver pharmaceuticals. For example, see US 5,401,516, US 5,443,841 and US RE35,862. In addition, certain modified
15 amino acids have been used to deliver pharmaceuticals. See, e.g., US 5,629,020; US 5,643,957; US 5,766,633; US 5,776,888; and US 5,866,536.

However, there is still a need for simple, inexpensive delivery systems which are easily prepared and which can
20 deliver a broad range of active agents by various routes.

SUMMARY OF THE INVENTION

Compounds and compositions that are useful in the
25 delivery of active agents are provided. The present invention encompasses compounds having the following formula, salts thereof, polymorphs thereof, or mixtures thereof.



Formula I

wherein

5 R^1 is C_1 - C_{10} alkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, C_3 - C_{10} cycloalkyl, phenyl, naphthyl, or aromatic heterocycle;

R^1 is optionally substituted with C_1 - C_4 alkyl or fluoroalkyl, C_2 - C_4 alkenyl, C_1 - C_4 alkoxy or fluoroalkoxy, halogens, -OH, -SH, phenyl, phenoxy, - CO_2R_3 , - $N(CH_3)_2$, - NO_2 , or
10 - NH_2 .

R^2 is C_1 - C_{24} alkylene, C_2 - C_{24} alkenylene, C_3 - C_{10} cycloalkylene, C_3 - C_{10} cycloalkenylene, phenylene, naphthylene, (C_1 - C_{10} alkyl)phenylene, (C_2 - C_{10} alkenyl)phenylene, (C_1 - C_{10} alkyl)naphthylene, (C_2 - C_{10} alkenyl)naphthylene, phenyl(C_1 - C_{10} alkylene), phenyl(C_2 - C_{10} alkenylene), naphthyl(C_1 - C_{10} alkylene) or naphthyl(C_2 - C_{10} alkenylene);

R^2 is optionally substituted with C_1 - C_4 alkyl or fluoroalkyl, C_2 - C_4 alkenyl, C_1 - C_4 alkoxy or fluoroalkoxy, halogens, -OH, -SH, phenyl, phenoxy, - CO_2R_3 , - $N(CH_3)_2$, - NO_2 , -
20 NH_2 , C_3 - C_{10} cycloalkyl, C_3 - C_{10} cycloalkenyl, aryl, (C_1 - C_{10} alk)aryl, heterocycle having 3-10 ring atoms wherein the hetero atom is one or more of N, O, S or any combination thereof, or any combination thereof;


R^2 being optionally interrupted by oxygen, nitrogen,
25 sulfur, or any combination thereof; and

R^3 is hydrogen, C_1 - C_4 alkyl, or C_2 - C_4 alkenyl, with the provisos that R^1 is not 4-(piperidin-4-yl)phenyl when R^2 is $-(CH_2)_4-$; R^1 is not $-CH_3$ when R^2 is $-(CH_2)_3-$; R^1 is not 4-carboxylphenyl when R^2 is $-(CH_2)_3-$ or $-(CH_2)_4-$.

5 Preferably, R^1 is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, cycloalkyl, phenyl, or naphthyl and R^1 is optionally substituted with halogens, $-OH$, $-SH$, C_1 - C_4 alkyl or C_1 - C_4 alkoxy. More preferably, R^1 is a substituted or unsubstituted phenyl or naphthyl, $-CH_3$, or $-CH_2OH$. Most preferably, R^1 is 2-
10 OH-phenyl, optionally further substituted with $-OH$, Cl , $-CH_3$ or $-OCH_3$.

Preferably, R^2 is C_1 - C_{24} alkylene or $(C_1$ - C_{10} alkyl)phenylene. More preferably, R^2 is C_4 - C_8 alkylene.


More preferred compounds include, but are not limited to,
15 those having the formulas:



wherein

Cpd #	X	R^2
1	2-OH	$(CH_2)_8$
2	2-OH	$(CH_2)_6$
3	2-OH	$(CH_2)_4$
4	2-OH	$(CH_2)_7$
5	2-OH-5-Cl	$(CH_2)_6$

Cpd #	X	R ²
6	2-OH-3,5-di- Cl	(CH ₂) ₆
7	2-OH-4- methyl	(CH ₂) ₆
8	2-OH-4-OMe	(CH ₂) ₆
9	2-OH-4-OMe	(CH ₂) ₄
10	H	(CH ₂) ₆
11	2-Cl	(CH ₂) ₆




wherein

Cpd#	R ¹	R ²
12	CH ₃	(CH ₂) ₆
13	CH ₂ OH	(CH ₂) ₆

polymorphs thereof, and salts thereof.

These compounds may be referred to as oxadiazoles.

The compositions of the present invention comprise at least one active agent, preferably a biologically or chemically active agent, and at least one of the compounds or salts thereof of the present invention of the structure of Formula I above, including all the particular embodiments referred to above as well as those excluded by proviso, and also including compounds 1-13. These compounds may have the formula



wherein

R¹ is C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₃-C₁₀ cycloalkyl, phenyl, naphthyl, or aromatic heterocycle;

5 R¹ is optionally substituted with C₁-C₄ alkyl or fluoroalkyl, C₂-C₄ alkenyl, C₁-C₄ alkoxy or fluoroalkoxy, halogens, -OH, -SH, phenyl, phenoxy, -CO₂R₃, -N(CH₃)₂, -NO₂, or -NH₂.

10 R² is C₁-C₂₄ alkylene, C₂-C₂₄ alkenylene, C₃-C₁₀ cycloalkylene, C₃-C₁₀ cycloalkenylene, phenylene, naphthylene, (C₁-C₁₀ alkyl)phenylene, (C₂-C₁₀ alkenyl)phenylene, (C₁-C₁₀ alkyl)naphthylene, (C₂-C₁₀ alkenyl)naphthylene, phenyl(C₁-C₁₀ alkylene), phenyl(C₂-C₁₀ alkenylene), naphthyl(C₁-C₁₀ alkylene) or naphthyl(C₂-C₁₀ alkenylene);

15 R² is optionally substituted with C₁-C₄ alkyl or fluoroalkyl, C₂-C₄ alkenyl, C₁-C₄ alkoxy or fluoroalkoxy, halogens, -OH, -SH, phenyl, phenoxy, -CO₂R₃, -N(CH₃)₂, -NO₂, -NH₂, C₃-C₁₀ cycloalkyl, C₃-C₁₀ cycloalkenyl, aryl, (C₁-C₁₀ alk)aryl, heterocycle having 3-10 ring atoms wherein the
20 hetero atom is one or more of N, O, S or any combination thereof, or any combination thereof;

R² being optionally interrupted by oxygen, nitrogen, sulfur, or any combination thereof; and

R³ is hydrogen, C₁-C₄ alkyl, or C₂-C₄ alkenyl.

25 Methods for the preparation and administration of such

compositions are also provided.

Also provided are dosage unit forms comprising the composition of the present invention. The dosing vehicle can be a solid (such as a tablet, powder, or capsule) or a liquid.

5 Methods for administering a biologically active agent to an animal in need of the agent, especially by the oral or intracolonic routes, with the compositions of the present invention, are also provided.

10

DETAILED DESCRIPTION OF THE INVENTION

Compounds

15 The compounds may be in the form of the carboxylic acid and/or their salts. Salts include but are not limited to organic and inorganic salts, for example alkali-metal salts, such as sodium, potassium and lithium; alkaline-earth metal salts, such as magnesium, calcium or barium; ammonium salts; basic amino acids, such as lysine or arginine; and organic
20 amines, such as dimethylamine or pyridine. Preferably, the salts are sodium salts.

 In general, the compounds of the present invention may be prepared by dehydrative cyclization of dihydrazides, protecting any functional groups present in the dihydrazide as
25 necessary. Dehydrative cyclization may be performed by reacting dihydrazide with carbon tetrachloride and triphenylphosphene in the presence of an organic base, such as triethylamine. The acid is formed from basic hydrolysis of the resulting ester. Suitable solvents for the cyclization
30 reaction include, but are not limited to, acetonitrile.

 The compound may be purified by recrystallization or by

fractionation on one or more solid chromatographic supports, alone or linked in tandem. Suitable recrystallization solvent systems include, but are not limited to, acetonitrile, methanol, and tetrahydrofuran. Fractionation may be performed
5 on a suitable chromatographic support such as alumina, using methanol/n-propanol mixtures as the mobile phase; reverse phase chromatography using trifluoroacetic acid/acetonitrile mixtures as the mobile phase; and ion exchange chromatography using water or an appropriate buffer as the mobile phase.
10 When anion exchange chromatography is performed, preferably a 0-500 mM sodium chloride gradient is employed.

Active Agents

Active agents suitable for use in the present invention
15 include biologically active agents and chemically active agents, including, but not limited to, pesticides, pharmacological agents, and therapeutic agents.

For example, biologically or chemically active agents suitable for use in the present invention include, but are not
20 limited to, proteins; polypeptides; peptides; hormones; polysaccharides, and particularly mixtures of mucopolysaccharides; carbohydrates; lipids; other organic compounds; and particularly compounds which by themselves do not pass (or which pass only a fraction of the administered
25 dose) through the gastro-intestinal mucosa and/or are susceptible to chemical cleavage by acids and enzymes in the gastro-intestinal tract; or any combination thereof.

Further examples include, but are not limited to, the following, including synthetic, natural or recombinant sources
30 thereof: growth hormones, including human growth hormones (hGH), recombinant human growth hormones (rhGH), bovine growth

hormones, and porcine growth hormones; growth hormone-releasing hormones; interferons, including α , β and γ ; interleukin-1; interleukin-2; insulin, including porcine, bovine, human, and human recombinant, optionally having
5 counter ions including sodium, zinc, calcium and ammonium; insulin-like growth factor, including IGF-1; heparin, including unfractionated heparin, heparinoids, dermatans, chondroitins, low molecular weight heparin, very low molecular weight heparin and ultra low molecular weight heparin;
10 calcitonin, including salmon, eel and human; erythropoietin; atrial natriuretic factor; antigens; monoclonal antibodies; somatostatin; protease inhibitors; adrenocorticotropin, gonadotropin releasing hormone; oxytocin; leutinizing-hormone-releasing-hormone; follicle stimulating hormone;
15 glucocerebrosidase; thrombopoietin; filgrastim; prostaglandins; cyclosporin; vasopressin; cromolyn sodium (sodium or disodium chromoglycate); vancomycin; desferrioxamine (DFO); parathyroid hormone (PTH), including its fragments; antimicrobials, including anti-fungal agents;
20 vitamins; analogs, fragments, mimetics or polyethylene glycol (PEG)-modified derivatives of these compounds; or any combination thereof.

Delivery systems

25 The compositions of the present invention comprise a delivery agent, i.e., a compound of the present invention, and one or more active agents. In one embodiment, one or more of the delivery agent compounds, or salts of these compounds, may be used as a delivery agent by mixing with the active agent
30 prior to administration.

The administration compositions may be in the form of a

liquid. The solution medium may be water (for example, for salmon calcitonin, parathyroid hormone, and erythropoietin), 25% aqueous propylene glycol (for example, for heparin) and phosphate buffer (for example, for rhGH). Other dosing
5 vehicles include polyethylene glycols, sorbitol, maltitol, and sucrose. Dosing solutions may be prepared by mixing a solution of the delivery agent compound with a solution of the active agent, just prior to administration. Alternately, a solution of the delivery agent (or active agent) may be mixed
10 with the solid form of the active agent (or delivery agent). The delivery agent compound and the active agent may also be mixed as dry powders. The delivery agent compound and the active agent can also be admixed during the manufacturing process.

15 The dosing solutions may optionally contain additives such as phosphate buffer salts, citric acid, glycols, or other dispersing agents. Stabilizing additives may be incorporated into the solution, preferably at a concentration ranging between about 0.1 and 20% (w/v).

20 The administration compositions may alternately be in the form of a solid, such as a tablet, capsule or powder. Solid dosage forms may be prepared by mixing the solid form of the compound with the solid form of the active agent. Alternately, a solid may be obtained from a solution of
25 compound and active agent by methods known in the art, such as freeze drying, precipitation, crystallization and solid dispersion.

The administration compositions of the present invention may also include one or more enzyme inhibitors. Such enzyme
30 inhibitors include, but are not limited to, compounds such as actinonin, epiactinonin, and derivatives thereof. Other

enzyme inhibitors include, but are not limited to, aprotinin (Trasylol) and Bowman-Birk inhibitor.

The amount of active agent used in an administration composition of the present invention is an amount effective to
5 accomplish the purpose of the particular active agent for the target indication. The amount of active agent in the compositions typically is a pharmacologically, biologically, therapeutically, or chemically effective amount. However, the amount can be less than that amount when the composition is
10 used in a dosage unit form because the dosage unit form may contain a plurality of compound/active agent compositions or may contain a divided pharmacologically, biologically, therapeutically, or chemically effective amount. The total effective amount can then be administered in cumulative units
15 containing, in total, an effective amount of the active agent.

The total amount of active agent to be used can be determined by methods known to those skilled in the art. However, because the compositions may deliver active agents more efficiently than prior compositions, lower amounts of
20 biologically or chemically active agents than those used in prior dosage unit forms or delivery systems can be administered to the subject, while still achieving the same blood levels and/or therapeutic effects.

The presently disclosed compounds facilitate delivery of
25 biologically and chemically active agents, particularly in oral, intranasal, sublingual, intraduodenal, subcutaneous, buccal, intracolonic, rectal, vaginal, mucosal, pulmonary, transdermal, intradermal, parenteral, intravenous, intramuscular and ocular systems, as well as traversing the
30 blood-brain barrier.

Dosage unit forms can also include any one or combination

of excipients, diluents, disintegrants, lubricants, plasticizers, colorants, flavorants, taste-masking agents, sugars, sweeteners, salts, and dosing vehicles, including, but not limited to, water, 1,2-propane diol, ethanol, olive oil,
5 or any combination thereof.

The compounds and compositions of the subject invention are useful for administering biologically or chemically active agents to any animals, including but not limited to birds such as chickens; mammals, such as rodents, cows, pigs, dogs, cats,
10 primates, and particularly humans; and insects.

The present invention is particularly advantageous for delivering chemically or biologically active agents that would otherwise be destroyed or rendered less effective by conditions encountered before the active agent reaches its
15 target zone (i.e. the area in which the active agent of the delivery composition is to be released) and within the body of the animal to which they are administered. Particularly, the compounds and compositions of the present invention are useful in orally administering active agents, especially those that
20 are not ordinarily orally deliverable, or those for which improved delivery is desired.

The compositions comprising the compounds and active agents have utility in the delivery of active agents to selected biological systems and in an increased or improved
25 bioavailability of the active agent compared to administration of the active agent without the delivery agent. Delivery can be improved by delivering more active agent over a period of time, or in delivering active agent in a particular time period (such as to effect quicker or delayed delivery) or over
30 a period of time (such as sustained delivery).

Following administration, the active agent present in the

composition or dosage unit form is taken up into the circulation. The bioavailability of the agent is readily assessed by measuring a known pharmacological activity in blood, e.g. an increase in blood clotting time caused by heparin, or a decrease in circulating calcium levels caused by calcitonin. Alternately, the circulating levels of the active agent itself can be measured directly.

10 DESCRIPTION OF THE PREFERRED EMBODIMENTS

The following examples illustrate the invention without limitation. All parts are given by weight unless otherwise indicated.

15 Example 1

Compound 2 was prepared as follows. A suspension of 30.34 g (0.108 mol) of N-(methyl suberoyl)-N'-salicyloylhydrazide and 138 mL of methylene chloride was cooled to 0 °C in an ice bath and treated with 16.56 mL (12.0 g, 0.119 mol) of triethylamine, followed by 14.40 mL (12.3 g, 0.113 mol) of trimethylsilylchloride 15 minutes later. After warming to 25 °C and stirring for 3 hours, the reaction mixture was concentrated in vacuo.

The residue was taken up in 183 mL of acetonitrile and treated with 45.16 mL (32.8 g, 0.324 mol) of triethylamine, 50.4 mL (80.2 g, 0.522 mol) of carbon tetrachloride and 63.88 g (0.243 mol) of triphenylphosphine. The orange slurry was stirred for 18 hours. The solid was removed by filtration. The filtrate was diluted with 183 mL of ethyl acetate, washed with 2% aqueous hydrochloric acid (2x100 mL) dried over sodium sulfate and concentrated in vacuo.

The resulting solid was brought up in 60 mL of 2N aqueous sodium hydroxide and stirred at 65 °C for three hours. The undissolved solid was removed by filtration. The filtrate was acidified to pH 3 with 3% aqueous hydrochloric acid. The
5 resulting solid was isolated by filtration and recrystallized from ethanol/water. A total of 11.76 g, having a melting point of 109-112 °C, was collected.

Compounds 1, 3-8, 10 and 12 were also prepared by this method with the appropriate starting materials. Compounds 9,
10 11 and 13 may also be prepared by this method using the appropriate starting materials.

Example 2 - Salmon Calcitonin (sCT)

Oral Delivery

15 Oral dosing (PO) compositions of delivery agent compound and salmon calcitonin (sCT) in water were prepared. Typically, 450 mg of compound was added to 2.0 ml of water. Either the sodium salt of the compound was used or the free acid was converted to the sodium salt by stirring the
20 resultant solution and adding one equivalent of sodium hydroxide (1.0 N) and diluting with water. The solution was vortexed, then heated (about 37 °C) and sonicated. The pH was adjusted to about 7 (6.5 to 8.5) with sodium hydroxide or hydrochloric acid. 90 µg sCT from a stock solution was added
25 to the solution. Water was then added to bring the total volume to about 3.0 ml (varies depending on solubility of the delivery agent compound) and vortexed. The final delivery agent compound dose, sCT dose and volume dose amounts are listed below in Table 1.

30 The typical dosing and sampling protocols were as

follows. Male Sprague-Dawley rats weighing between 200-250g were fasted for 24 hours and administered ketamine (44 mg/kg) and chlorpromazine (1.5 mg/kg) 15 minutes prior to dosing. A dosing group of five rats was administered one of the dosing solutions. For oral dosing, an 11 cm Rusch 8 French catheter was adapted to a 1 ml syringe with a pipette tip. The syringe was filled with dosing solution by drawing the solution through the catheter, which was then wiped dry. The catheter was placed down the esophagus leaving 1 cm of tubing past the incisors. Solution was administered by pressing the syringe plunger.

Blood samples were collected serially from the tail artery, typically at 0, 10, 20, 30, 60 and 90 minutes after administration. Serum sCT was determined by testing with an EIA kit (Kit # EIAS-6003 from Peninsula Laboratories, Inc., San Carlos, CA). Numbers were adjusted according to baseline values obtained at 0 minutes. The limit of the assay is 80 pg/ml. The results from the five rats in each dosing group were averaged for each time point. The maximum is reported below in Table 1.

Table 1
Oral Delivery of Salmon Calcitonin (sCT)

Compound	Volume Dose (ml/kg)	Compound Dose (mg/kg)	sCT Dose (µg/kg)	Mean Peak Serum Sct (pg/ml ± SD) (SE)
1	1	150	30	167 ± 89 (39)
2	1	150	30	260 ± 146 (65)
2	1	150	30	112 ± 69 (31)
2	1	150	30	165 ± 136 (61)
2	1	150	30	160 ± 207 (93)
2	1	150	30	492 ± 830 (10)
*3	1	150	30	271 ± 503 (225)
3	1	150	30	444 ± 277 (124)
3	1	150	30	108 ± 115 (52)
4	1	150	30	272 ± 284 (127)
4	1	150	30	253 ± 374 (167)
5	1	150	30	393 ± 431 (193)
6	1	150	30	306 ± 339 (152)
8	1	150	30	168 ± 217 (97)
8	1	150	30	204 ± 227 (102)
10	1	150	30	37 ± 55
12	1	150	30	212 ± 194 (433)
12	1	150	30	155 ± 134 (60)

* In this dosing of compound 3, the standard protocol was modified from the kit as follows: incubated with 50 µl peptide antibody for 2 hours with shaking in the dark, washed the plate, added serum and biotinylated peptide and diluted with 4 ml buffer, and shook overnight in the dark.

10

Example 3 - Heparin Delivery

Oral/Intracolonic Delivery

Oral gavage (PO) and/or intracolonic (IC) dosing solutions containing a delivery agent compound and heparin sodium USP in 25% aqueous propylene glycol were prepared.

15

Either the sodium salt of the compound was used or the free acid was converted to the sodium salt with one equivalent of sodium hydroxide. Typically, delivery agent compound and heparin (about 166-182 IU/mg) were mixed by vortex as dry
5 powders. This dry mixture was dissolved in 25% v/v aqueous propylene glycol, vortexed and placed in a sonicator (about 37 °C). The pH was adjusted to about 7 (6.5 to 8.5) with aqueous sodium hydroxide (2N). The dosing solution was sonicated to produce a clear solution. The final volume was adjusted to
10 about 3.0 ml. The final delivery agent compound dose, heparin dose and volume dose amounts are listed below in Table 2.

The typical dosing and sampling protocols were as follows. Male Sprague-Dawley rats weighing between 275-350g were fasted for 24 hours and were anesthetized with ketamine
15 hydrochloride (88 mg/kg) intramuscularly immediately prior to dosing. A dosing group of five rats was administered one of the dosing solutions. For oral gavage (PO) dosing, an 11cm Rusch 8 French catheter was adapted to a 1 ml syringe with a pipette tip. The syringe was filled with dosing solution by
20 drawing the solution through the catheter, which was then wiped dry. The catheter was placed down the esophagus leaving 1 cm of tubing past the rat's incisors. Solution was administered by pressing the syringe plunger. For
25 intracolonic (IC) dosing, a 7.5cm, 8 fr Rusch catheter was adapted to a 1 ml syringe with a pipette tip. The dosing catheter was inserted into the colon through the anus until the tube was no longer visible. The dosing solution was expressed slowly into the colon by pressing the syringe
30 plunger.

Citrated blood samples were collected by cardiac puncture following the administration of ketamine (88 mg/kg), typically

at 0.25, 0.5, 1.0 and 1.5 hours after administration. Heparin activity was determined by utilizing the activated partial thromboplastin time (APTT) according to the method of Henry, J.B., Clinical Diagnosis and Management by Laboratory Methods, Philadelphia, PA, W.B. Saunders (1979). Previous studies indicated baseline values of about 20 sec. Results from the five rats in each group were averaged for each time point. The maximum is reported below in Table 2.

Table 2

Oral/Intracolonic Delivery of Heparin

Compound	Method of Administration	Volume dose (ml/kg)	Compound Dose (mg/kg)	Heparin Dose (mg/kg)	Mean Peak APTT (sec) \pm SD
1	IC	1	50	25	38.4 \pm 17.4
2	IC	1	50	25	52.8 \pm 31.9

Example 4 - Recombinant Human Growth Hormone (rhGH)

Oral/Intracolonic Delivery

Oral gavage (PO) and/or intracolonic (IC) dosing solutions of delivery agent compound and rhGH in phosphate buffer were prepared. A solution of the compound was made either with the sodium salt of the compound or by converting the free acid to its sodium salt. Typically, a solution of the compound was prepared in phosphate buffer and stirred, adding one equivalent of sodium hydroxide (1.0 N) when making the sodium salt. The final dosing solutions were prepared by mixing the compound solution with an rhGH stock solution (15 mg rhGH/ml) and diluting to the desired volume (usually 3.0 ml). The compounds and rhGH dose amounts are listed below in

Table 3.

The typical dosing and sampling protocols were as follows. Male Sprague-Dawley rats weighing between 200-250g were fasted for 24 hours and administered ketamine (44 mg/kg) and chlorpromazine (1.5 mg/kg) 15 minutes prior to dosing. A dosing group of five rats was administered one of the dosing solutions. For oral gavage (PO), an 11cm Rusch 8 French catheter was adapted to a 1 ml syringe with a pipette tip. The syringe was filled with dosing solution by drawing the solution through the catheter, which was then wiped dry. The catheter was placed down the esophagus leaving 1 cm of tubing past the rat's incisors. Solution was administered by pressing the syringe plunger. For intracolonic (IC) dosing, a 7.5 cm Rusch catheter tube (French 8 or 6) was adapted to a syringe with an Eppendorf pipette tip. The syringe was filled with the dosing solution by drawing the solution through the catheter tube. The catheter tube was wiped dry. K-Y jelly was applied to the tip, avoiding contact with the eye of the tube, and the tube was inserted into the colon through the anus until the tube was no longer visible. The solution was injected by pressing the syringe plunger, and the tube was removed.

Blood samples were collected serially from the tail artery, typically at 0, 15, 30, 45, 60 and 90 minutes after oral administration and 0, 10, 20, 30, 60 and 90 minutes after IC administration. The five samples from each time period were pooled (except for where compound 1 was dosed). Serum rHGH concentrations were quantified by an rHGH immunoassay test kit (Kit # K1F4015 from Genzyme Corporation Inc., Cambridge, MA). Previous studies indicated baseline values of about zero.

The maximum concentration for each group and the area under the curve (AUC), when available, are reported below in Table 3.

5

Table 3

Oral/Intracolonic Delivery of rhGH in Rats

Compound	Method of Administration	Volume dose (ml/kg)	Compound Dose (mg/kg)	rhGH Dose (mg/kg)	Mean Peak Serum [rhGH] (ng/ml)	AUC
1	PO	1	200	3	4.42 ± 4.42	142.25
1	PO	1	200	3	11.1 ± 8.9	218.7
2	PO	1	200	3	3.77	56.55
3	PO	1	200	3	88.9	133.5
3	PO	1	200	3	35.08	969.6
5	PO	1	200	3	14.64	219.6
6	PO	1	200	3	0	0
7	PO	1	200	3	0.045	60
8	PO	1	200	3	9.81	2187.7
8	PO	1	200	3	0	0
9	PO	1	200	3	14.25	
10	PO	1	200	3	14.25	802.725
12	PO	1	200	3	0	0

10 Example 5 - Parathyroid Hormone Delivery (PTH 1-34)Oral/Intracolonic Delivery

Oral gavage (PO) and/or intracolonic (IC) dosing solutions of delivery agent compound and human parathyroid hormone residues 1-34 (PTH) in water were prepared. A solution of the compound was made either with the sodium salt of the compound or by converting the free acid to its sodium salt. Typically, a solution of the compound was prepared in water and stirred, adding one equivalent of sodium hydroxide (1.0 N) when making the sodium salt. The final dosing solutions were prepared by mixing the compound solution with a

20

PTH stock solution (typically having a concentration of 5 mg PTH/ml) and diluting to the desired volume (usually 3.0 ml). The final compound, PTH and volume dose amounts are listed below in Table 4.

5 The typical dosing and sampling protocols were as follows. Male Sprague-Dawley rats weighing between 200-250g were fasted for 24 hours and administered ketamine (44 mg/kg) and chlorpromazine (1.5 mg/kg) 15 minutes prior to dosing. A dosing group of five rats was administered one of the dosing
10 solutions. For oral gavage (PO), an 11cm Rusch 8 French catheter was adapted to a 1 ml syringe with a pipette tip. The syringe was filled with dosing solution by drawing the solution through the catheter, which was then wiped dry. The catheter was placed down the esophagus leaving 1 cm of tubing
15 past the rat's incisors. Solution was administered by pressing the syringe plunger. For intracolonic (IC) dosing, a 7.5 cm Rusch catheter tube (French 8 or 6) was adapted to a syringe with an Eppendorf pipette tip. The syringe was filled with the dosing solution by drawing the solution through the
20 catheter tube. The catheter tube was wiped dry. K-Y jelly was applied to the tip, avoiding contact with the eye of the tube, and the tube was inserted into the colon through the anus until the tube was no longer visible. The solution was injected by pressing the syringe plunger, and the tube was
25 removed.

 Blood samples were collected serially from the tail artery, typically 0, 15, 30, 45, 60 and 90 minutes after oral administration and 0, 10, 20, 30, 60 and 90 minutes after IC administration. Serum PTH concentrations were quantified by a
30 PTH radioimmunoassay kit (Kit # RIK 6101 from Peninsula Laboratories, Inc., San Carlos, CA). Previous studies

indicated baseline values of about zero. Results from the five rats in each group were averaged for each time point. The maximum is reported below in Table 4.

5

Table 4
Oral/Intracolonic Delivery of PTH in Rats


Compound	Method of Administration	Volume dose (ml/kg)	Compound Dose (mg/kg)	PTH Dose (μ g/kg)	Mean Peak Serum [PTH] (pg/ml) \pm SD	AUC
2	PO	1	100	200	498.55 \pm 237.83	
3	PO	1	100	200	57.88 \pm	2717.51
4	PO	1	100	200	330.74 \pm 410.57	6852.12
5	PO	1	100	200	22.91 \pm	509.20
6	PO	1	100	200	14.44 \pm	716.46
7	PO	1	100	200	138.9 \pm 118.46	4790.10

10 The above mentioned patents, applications, test methods, and publications are hereby incorporated by reference in their entirety.


15 Many variations of the present invention will suggest themselves to those skilled in the art in light of the above detailed description. All such obvious variations are within the fully intended scope of the appended claims

WHAT IS CLAIMED IS:

1. A compound selected from the group consisting of



Cpd #	X	R ²
1	2-OH	(CH ₂) ₈
2	2-OH	(CH ₂) ₆
3	2-OH	(CH ₂) ₄
4	2-OH	(CH ₂) ₇
5	2-OH-5-Cl	(CH ₂) ₆
6	2-OH-3,5-di- Cl	(CH ₂) ₆
7	2-OH-4- methyl	(CH ₂) ₆
8	2-OH-4-OMe	(CH ₂) ₆
9	2-OH-4-OMe	(CH ₂) ₄
10	H	(CH ₂) ₆
11	2-Cl	(CH ₂) ₆



10

Cpd#	R ¹	R ²
12	CH ₃	(CH ₂) ₆
13	CH ₂ OH	(CH ₂) ₆

11

12 and salts thereof.

1 2. A composition comprising:

2 (A) an active agent; and

3 (B) at least one compound of claim 1.

1 3. The composition of claim 2, wherein the active agent is
2 selected from the group consisting of a biologically active
3 agent, a chemically active agent, and a combination thereof.

1 4. The composition of claim 3, wherein the biologically
2 active agent comprises at least one protein, polypeptide,
3 peptide, hormone, polysaccharide, mucopolysaccharide,
4 carbohydrate, or lipid.


1 5. The composition of claim 3, wherein the biologically
2 active agent is selected from the group consisting of growth
3 hormones, human growth hormones (hGH), recombinant human growth
4 hormones (rhGH), bovine growth hormones, porcine growth hormones,
5 growth hormone-releasing hormones, interferons, α -interferon, β -
6 interferon, γ -interferon, interleukin-1, interleukin-2, insulin,
7 porcine insulin, bovine insulin, human insulin, human recombinant
8 insulin, insulin-like growth factor(IGF), IGF-1, heparin,
9 unfractionated heparin, heparinoids, dermatans, chondroitins, low
10 molecular weight heparin, very low molecular weight heparin,
11 ultra low molecular weight heparin, calcitonin, salmon
12 calcitonin, eel calcitonin, human calcitonin; erythropoietin

13 (EPO), atrial naturetic factor, antigens, monoclonal antibodies,
14 somatostatin, protease inhibitors, adrenocorticotropin,
15 gonadotropin releasing hormone, oxytocin, leutinizing-hormone-
16 releasing-hormone, follicle stimulating hormone,
17 glucocerebrosidase, thrombopoietin, filgrastim, prostaglandins,
18 cyclosporin, vasopressin, cromolyn sodium, sodium chromoglycate,
19 disodium chromoglycate, vancomycin, desferrioxamine (DFO),
20 parathyroid hormone (PTH), fragments of PTH, antimicrobials,
21 anti-fungal agents, vitamins; analogs, fragments, mimetics and
22 polyethylene glycol (PEG)-modified derivatives of these
23 compounds; and any combination thereof.

1 6. The composition of claim 3, wherein the biologically
2 active agent comprises insulin, unfractionated heparin, low
3 molecular weight heparin, very low molecular weight heparin,
4 ultra low molecular weight heparin, calcitonin, PTH, EPO, hGH or
5 combinations thereof.

1 7. The composition of claim 3, wherein the biologically
2 active agent comprises salmon calcitonin.

1 8. The composition of claim 2, wherein the compound is
2



3
4 wherein

Cpd #	X	R ²
3	2-OH	(CH ₂) ₄

5
6 or a salt thereof.

8 9. A dosage unit form comprising:
9 (A) the composition of claim 2; and
10 (B) (a) an excipient
11 (b) a diluent,
12 (c) a disintegrant,
13 (d) a lubricant,
14 (e) a plasticizer,
15 (f) a colorant,
16 (g) a dosing vehicle, or
17 (h) any combination thereof.

1 10. The dosage unit form of claim 9, wherein the
2 biologically active agent comprises at least one protein,
3 polypeptide, peptide, hormone, polysaccharide,
4 mucopolysaccharide, carbohydrate, or lipid.
5

1 11. The dosage unit form of claim 9, wherein the
2 biologically active agent is selected from the group consisting
3 of growth hormones, human growth hormones (hGH), recombinant
4 human growth hormones (rhGH), bovine growth hormones, porcine
5 growth hormones, growth hormone-releasing hormones, interferons,
6 α -interferon, β -interferon, γ -interferon, interleukin-1,
7 interleukin-2, insulin, porcine insulin, bovine insulin, human
8 insulin, human recombinant insulin, insulin-like growth
9 factor(IGF), IGF-1, heparin, unfractionated heparin, heparinoids,
10 dermatans, chondroitins, low molecular weight heparin, very low
11 molecular weight heparin, ultra low molecular weight heparin,
12 calcitonin, salmon calcitonin, eel calcitonin, human calcitonin;
13 erythropoietin (EPO), atrial natriuretic factor, antigens,
14 monoclonal antibodies, somatostatin, protease inhibitors,
15 adrenocorticotropin, gonadotropin releasing hormone, oxytocin,
16 leutinizing-hormone-releasing-hormone, follicle stimulating
17 hormone, glucocerebrosidase, thrombopoietin, filgrastim,
18 prostaglandins, cyclosporin, vasopressin, cromolyn sodium, sodium
19 chromoglycate, disodium chromoglycate, vancomycin,
20 desferrioxamine (DFO), parathyroid hormone (PTH), fragments of
21 PTH, antimicrobials, anti-fungal agents, vitamins; analogs,
22 fragments, mimetics and polyethylene glycol (PEG)-modified
23 derivatives of these compounds; and any combination thereof.

25 12. The dosage unit form of claim 10, wherein the
26 biologically active agent comprises insulin, unfractionated
27 heparin, low molecular weight heparin, very low molecular weight
28 heparin, ultra low molecular weight heparin, calcitonin, PTH,
29 EPO, hGH, or combinations thereof.


1 13. The dosage unit form of claim 9, wherein the active
2 agent comprises salmon calcitonin.

1 14. The dosage unit form of claim 9, wherein the dosage
2 unit form is in the form of a tablet, a capsule, a powder, or a
3 liquid.

1 15. A method for administering a biologically-active agent
2 to an animal in need of the agent, the method comprising
3 administering orally to the animal the composition of claim 3.

1 16. A method for preparing a composition comprising
2 mixing:
3 (A) at least one active agent;
4 (B) the compound of claim 1; and
5 (C) optionally, a dosing vehicle.

1 17. A compound having the formula



2

3 wherein

4 R^1 is C_1 - C_{10} alkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, C_3 - C_{10}
5 cycloalkyl, phenyl, naphthyl, or aromatic heterocycle;

6 R^1 is optionally substituted with C_1 - C_4 alkyl or fluoroalkyl,
7 C_2 - C_4 alkenyl, C_1 - C_4 alkoxy or fluoroalkoxy, halogens, -OH, -SH,
8 phenyl, phenoxy, $-CO_2R_3$, $-N(CH_3)_2$, $-NO_2$, or $-NH_2$.

9 R^2 is C_1 - C_{24} alkylene, C_2 - C_{24} alkenylene, C_3 - C_{10} cycloalkylene,
10 C_3 - C_{10} cycloalkenylene, phenylene, naphthylene, (C_1 - C_{10}
11 alkyl)phenylene, (C_2 - C_{10} alkenyl)phenylene, (C_1 - C_{10}
12 alkyl)naphthylene, (C_2 - C_{10} alkenyl)naphthylene, phenyl(C_1 - C_{10}
13 alkylene), phenyl(C_2 - C_{10} alkenylene), naphthyl(C_1 - C_{10} alkylene) or
14 naphthyl(C_2 - C_{10} alkenylene);

15 R^2 is optionally substituted with C_1 - C_4 alkyl or fluoroalkyl,
16 C_2 - C_4 alkenyl, C_1 - C_4 alkoxy or fluoroalkoxy, halogens, -OH, -SH,
17 phenyl, phenoxy, $-CO_2R_3$, $-N(CH_3)_2$, $-NO_2$, $-NH_2$, C_3 - C_{10} cycloalkyl,
18 C_3 - C_{10} cycloalkenyl, aryl, (C_1 - C_{10} alk)aryl, heterocycle having 3-
19 10 ring atoms wherein the hetero atom is one or more of N, O, S
20 or any combination thereof, or any combination thereof;


21 R^2 being optionally interrupted by oxygen, nitrogen, sulfur,
22 or any combination thereof; and

23 R^3 is hydrogen, C_1 - C_4 alkyl, or C_2 - C_4 alkenyl,
24 with the provisos that R^1 is not 4-(piperidin-4-yl)phenyl when R^2
25 is $-(CH_2)_4-$; R^1 is not $-CH_3$ when R^2 is $-(CH_2)_3-$; R^1 is not 4-
26 carboxylphenyl when R^2 is $-(CH_2)_3-$ or $-(CH_2)_4-$.

1 18. The compound of claim 17, wherein
 2 R^1 is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, or
 3 cycloalkyl, or phenyl or naphthyl, optionally substituted with
 4 halogens, -OH, -SH, C_1 - C_4 alkyl or C_1 - C_4 alkoxy; and
 5 R^2 is C_1 - C_{24} alkylene or (C_1 - C_{10} alkyl)phenylene.

1 19. The compound of claim 18, wherein
 2 R^1 is 2-OH-phenyl, optionally further substituted with
 3 -OH, Cl, -CH₃ or -OCH₃;
 4 R^2 is C_4 - C_8 alkyl.

1 20. A composition comprising:
 2 (A) an active agent; and
 3 (B) a compound having the formula



4
 5 wherein
 6 R^1 is C_1 - C_{10} alkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, C_3 - C_{10}
 7 cycloalkyl, phenyl, naphthyl, or aromatic heterocycle;
 8 R^1 is optionally substituted with C_1 - C_4 alkyl or fluoroalkyl,
 9 C_2 - C_4 alkenyl, C_1 - C_4 alkoxy or fluoroalkoxy, halogens, -OH, -SH,
 10 phenyl, phenoxy, -CO₂R₃, -N(CH₃)₂, -NO₂, or -NH₂.
 11 R^2 is C_1 - C_{24} alkylene, C_2 - C_{24} alkenylene, C_3 - C_{10} cycloalkylene,
 12 C_3 - C_{10} cycloalkenylene, phenylene, naphthylene, (C_1 - C_{10}
 13 alkyl)phenylene, (C_2 - C_{10} alkenyl)phenylene, (C_1 - C_{10}
 14 alkyl)naphthylene, (C_2 - C_{10} alkenyl)naphthylene, phenyl(C_1 - C_{10}
 15 alkylene), phenyl(C_2 - C_{10} alkenylene), naphthyl(C_1 - C_{10} alkylene) or

17 naphthyl(C₂-C₁₀ alkenylene);

18 R² is optionally substituted with C₁-C₄ alkyl or fluoroalkyl,
19 C₂-C₄ alkenyl, C₁-C₄ alkoxy or fluoroalkoxy, halogens, -OH, -SH,
20 phenyl, phenoxy, -CO₂R₃, -N(CH₃)₂, -NO₂, -NH₂, C₃-C₁₀ cycloalkyl,
21 C₃-C₁₀ cycloalkenyl, aryl, (C₁-C₁₀ alk)aryl, heterocycle having 3-
22 10 ring atoms wherein the hetero atom is one or more of N, O, S
23 or any combination thereof, or any combination thereof;

24 R² being optionally interrupted by oxygen, nitrogen, sulfur,
25 or any combination thereof; and

26 R³ is hydrogen, C₁-C₄ alkyl, or C₂-C₄ alkenyl.

INTERNATIONAL SEARCH REPORT

 International application No.
PCT/US00/03899

A. CLASSIFICATION OF SUBJECT MATTER IPC(7) : A61K 9/20, 9/48, 31/415 US CL : 424/451,464; 514/403 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 424/451,464; 514/403 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) WEST, CAPLUS, CAOLD, BIOSIS, EMBASE, MEDLINE		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim 1
Y	ANDOTRA, C.S. Synthesis and biocidal activity of some 1, 3, 4-oxadiazolo(3,2-a)-s-triazine-5-7-dithiones. Indian J. Heterocycl. Chem. June 1991, Vol. 1, No. 1, pages 26-28, see entire document.	1-20
Y	KAHN et al. Insulin inhibits vascular smooth muscle contraction at a site distal to intracellular Ca ²⁺ concentration. Am. J. Physiol. 1998, Vol. 274, No. 37, pages E885-E892, see entire document.	1-20
Y	HASDAI et al. Insulin and Insulin-like Growth Factor-I Cause Coronary Vasorelaxation In Vitro. Hypertension. August 1998, Vol. 32, pages 228-234, see entire document.	1-20
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: *A* document defining the general state of the art which is not considered to be of particular relevance *B* earlier document published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art *&* document member of the same patent family	
Date of the actual completion of the international search 02 MAY 2000		Date of mailing of the international search report 05 JUL 2000
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer JOSEPH F. MURPHY Telephone No. (703) 308-0196